

Due Sun

nonempty set w/ vector add, scalar mult.
satisfying 10 axioms

6.1 - Inner Products

Definition: An inner product on a real vector space V is a function that associates a real number $\langle \mathbf{u}, \mathbf{v} \rangle$ with each pair of vectors in V in such a way that the following axioms are satisfied for all vectors \mathbf{u}, \mathbf{v} , and \mathbf{w} in V and all scalars k .

1. $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$ (symmetry axiom)
2. $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle + \langle \mathbf{v}, \mathbf{w} \rangle$ (additivity axiom)
3. $\langle k\mathbf{u}, \mathbf{v} \rangle = k \langle \mathbf{u}, \mathbf{v} \rangle$ (homogeneity axiom)
4. $\langle \mathbf{v}, \mathbf{v} \rangle \geq 0$ and $\langle \mathbf{v}, \mathbf{v} \rangle = 0$ if and only if $\mathbf{v} = \mathbf{0}$ (positivity axiom)

Definition: A real vector space with an inner product is called a **real inner product space**.

Note that an inner product is a category of operation that is performed on vectors in a vector space. Any operation that satisfies these axioms is an inner product.

collection of objects \rightarrow set

set together with
particularly defined
addition and scalar mult \rightarrow vector space

vector space with an
inner product \rightarrow inner product
space

Examples of Inner products

- The dot product $\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + \dots + u_nv_n$ is the **Euclidean inner product** or **standard inner product** on R^n . [R^n with the Euclidean inner product is called **Euclidean n-space**.]
- If $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$ are vectors in R^n and w_1, w_2, \dots, w_n are positive real numbers, then the formula $\langle \mathbf{u}, \mathbf{v} \rangle = w_1u_1v_1 + w_2u_2v_2 + \dots + w_nu_nv_n$ is called the **weighted Euclidean inner product with weights w_1, w_2, \dots, w_n** .
- On M_{nn} , the set of $n \times n$ matrices: If $\mathbf{u} = U$ and $\mathbf{v} = V$ are matrices in the vector space M_{nn} , then the formula $\langle \mathbf{u}, \mathbf{v} \rangle = \text{tr}(U^T V)$ is the **standard inner product** on M_{nn} .
- If $\mathbf{p} = a_0 + a_1x + \dots + a_nx^n$ and $\mathbf{q} = b_0 + b_1x + \dots + b_nx^n$ are polynomials in P_n , then the **standard inner product** on P_n is $\langle \mathbf{p}, \mathbf{q} \rangle = a_0b_0 + a_1b_1 + \dots + a_nb_n$. (Note the similarity in form to the dot product.)
- If $\mathbf{p} = a_0 + a_1x + \dots + a_nx^n$ and $\mathbf{q} = b_0 + b_1x + \dots + b_nx^n$ are polynomials in P_n and if x_0, x_1, \dots, x_n are distinct real numbers, then the formula $\langle \mathbf{p}, \mathbf{q} \rangle = p(x_0)q(x_0) + p(x_1)q(x_1) + \dots + p(x_n)q(x_n)$ is the **evaluation inner product** at x_0, x_1, \dots, x_n .
- If $\mathbf{f} = f(x)$ and $\mathbf{g} = g(x)$ are two functions in $C[a, b]$, then $\langle \mathbf{f}, \mathbf{g} \rangle = \int_a^b f(x)g(x)dx$ defines an **inner product** on $C[a, b]$.
- If \mathbf{u} and \mathbf{v} are vectors in R^n expressed in column form, A is an invertible $n \times n$ matrix, and $\mathbf{u} \cdot \mathbf{v}$ is the Euclidean inner product on R^n , then the formula $\langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{A}\mathbf{u} \cdot \mathbf{A}\mathbf{v}$ is the **inner product on R^n generated by A** . This is an example of a **matrix inner product**.

#10 Compute the standard inner product on M_{22} of the given matrices.

$$U = \begin{bmatrix} 1 & 2 \\ -3 & 5 \end{bmatrix}, V = \begin{bmatrix} 4 & 6 \\ 0 & 8 \end{bmatrix} \quad \langle \vec{u}, \vec{v} \rangle = \text{tr}(U^T V)$$

$$U^T = \begin{bmatrix} 1 & -3 \\ 2 & 5 \end{bmatrix} \Rightarrow U^T V = \begin{bmatrix} 1 & -3 \\ 2 & 5 \end{bmatrix} \begin{bmatrix} 4 & 6 \\ 0 & 8 \end{bmatrix}$$

$$U^T V = \begin{bmatrix} 4 & -18 \\ 8 & 52 \end{bmatrix} \Rightarrow \text{tr}(U^T V) = 4 + 52 = \boxed{56}$$

Consider $U = \begin{bmatrix} u_1 & u_2 \\ u_3 & u_4 \end{bmatrix}, V = \begin{bmatrix} v_1 & v_2 \\ v_3 & v_4 \end{bmatrix}$

Then $U^T V = \begin{bmatrix} u_1 & u_3 \\ u_2 & u_4 \end{bmatrix} \begin{bmatrix} v_1 & v_2 \\ v_3 & v_4 \end{bmatrix} = \begin{bmatrix} u_1 v_1 + u_3 v_3 & u_1 v_2 + u_3 v_4 \\ v_1 u_2 + v_3 u_4 & u_2 v_2 + u_4 v_4 \end{bmatrix}$

$$\text{tr}(U^T V) = u_1 v_1 + u_2 v_2 + u_3 v_3 + u_4 v_4$$

#11 Find the standard inner product on P_2 of the given polynomials.

$$p = -2 + x + 3x^2, q = 4 - 7x^2$$

For $\vec{p} = a_0 + a_1 x + a_2 x^2, q = b_0 + b_1 x + b_2 x^2,$

$$\langle \vec{p}, \vec{q} \rangle = a_0 b_0 + a_1 b_1 + a_2 b_2$$

here, $\langle \vec{p}, \vec{q} \rangle = -2(4) + 1(0) + 3(-7) = \boxed{-29}$

#16 A sequence of sample points is given. Use the evaluation inner product on P_3 at those sample points to find $\langle p, q \rangle$ for the polynomials $p = x + x^3$ and $q = 1 + x^2$.
 $x_0 = -1, x_1 = 0, x_2 = 1, x_3 = 2$

$$\langle \vec{p}, \vec{q} \rangle = (-2)(2) + (0)(1) + (2)(2) + (10)(5) = 50$$

x	$p(x)$	$q(x)$
-1	-2	2
0	0	1
1	2	2
2	10	5

#8 Use the inner product on R^2 generated by the matrix A to find $\langle u, v \rangle$ for the vectors

$u = (0, -3)$ and $v = (6, 2)$.

$$\langle \vec{u}, \vec{v} \rangle = A\vec{u} \cdot A\vec{v}$$

$$A = \begin{bmatrix} 2 & 1 \\ -1 & 3 \end{bmatrix}$$

$$A\vec{u} = \begin{bmatrix} 2 & 1 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 0 \\ -3 \end{bmatrix} = \begin{bmatrix} -3 \\ -9 \end{bmatrix}, \quad A\vec{v} = \begin{bmatrix} 2 & 1 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 6 \\ 2 \end{bmatrix} = \begin{bmatrix} 14 \\ 0 \end{bmatrix}$$

$$A\vec{u} \cdot A\vec{v} = (-3, -9) \cdot (14, 0) = -42$$

Since $\vec{u} \cdot \vec{v} = \vec{v}^T \vec{u}$ (See Table 1.2), we have

$$\langle \vec{u}, \vec{v} \rangle = (A\vec{v})^T A\vec{u} = \vec{v}^T A^T A \vec{u}$$

Example: A weighted Euclidean inner product

For this class, we have HW: 10%, Exams: 60%, Final: 30%. Suppose we have 4 exams, each with 100 pts possible. The final has 100 pts possible, and total possible HW pts is 320.

Scores for Student X	Total possible points
HW: 265	320
Exams: 350	400
Final: 85	100

$\vec{u} = (265, 350, 85)$ is points earned

$\vec{v} = \left(\frac{1}{320}, \frac{1}{400}, \frac{1}{100}\right)$ is the base for each category

$\langle \vec{u}, \vec{v} \rangle = \vec{u} \cdot \vec{v}$ does not factor category weights

Define $\langle \vec{u}, \vec{v} \rangle = 10u_1v_1 + 60u_2v_2 + 30u_3v_3$.

$$\text{Then } \langle \vec{u}, \vec{v} \rangle = 10\left(\frac{265}{320}\right) + 60\left(\frac{350}{400}\right) + 30\left(\frac{85}{100}\right) \approx 86$$

This is a weighted Euclidean inner product with weights $w_1 = 10$, $w_2 = 60$, $w_3 = 30$.

Considering a matrix inner product, a weighted Euclidean inner product is generated by

$$A = \begin{bmatrix} \sqrt{w_1} & & 0 \\ & \sqrt{w_2} & \\ 0 & & \dots & \sqrt{w_n} \end{bmatrix} \text{ so that } A^T A = \begin{bmatrix} w_1 & & 0 \\ & w_2 & \\ 0 & & \dots & w_n \end{bmatrix}$$

Definition: If V is a real inner product space, then the **norm** or **length** of a vector \mathbf{v} in V is denoted by $\|\mathbf{v}\|$ and is defined by $\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}$ and the **distance** between two vectors is denoted by $d(\mathbf{u}, \mathbf{v})$ and is defined by $d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\| = \sqrt{\langle \mathbf{u} - \mathbf{v}, \mathbf{u} - \mathbf{v} \rangle}$. A vector of norm 1 is called a **unit vector**.

#1 Let R^2 have the weighted Euclidean inner product $\langle \mathbf{u}, \mathbf{v} \rangle = 2u_1v_1 + 3u_2v_2$ and let $\mathbf{u} = (1, 1)$, $\mathbf{v} = (3, 2)$, $\mathbf{w} = (0, -1)$, and $k = 3$. Compute the stated quantities.
 a. $\langle \mathbf{u}, \mathbf{v} \rangle$ b. $\langle k\mathbf{u}, \mathbf{v} \rangle$ c. $\langle \mathbf{u} + \mathbf{v}, \mathbf{w} \rangle$ d. $\|\mathbf{v}\|$ e. $d(\mathbf{u}, \mathbf{v})$ f. $\|\mathbf{u} - k\mathbf{v}\|$

$$a) \langle \vec{u}, \vec{v} \rangle = 2(1)(3) + 3(1)(2) = 12$$

$$b) \langle 3\vec{u}, \vec{v} \rangle = 2(3)(3) + 3(3)(2) = 36 \quad \text{Note: } \langle 3\vec{u}, \vec{v} \rangle = 3\langle \vec{u}, \vec{v} \rangle$$

$$c) \langle \vec{u} + \vec{v}, \vec{w} \rangle = 2(1+0)(0) + 3(1+2)(-1) = -9$$

$$\text{Note: } \langle \vec{u}, \vec{w} \rangle = -3, \langle \vec{v}, \vec{w} \rangle = -6$$

$$\langle \vec{u}, \vec{w} \rangle + \langle \vec{v}, \vec{w} \rangle = \langle \vec{u} + \vec{v}, \vec{w} \rangle$$

$$d) \|\vec{v}\| = \sqrt{\langle \vec{v}, \vec{v} \rangle} = \sqrt{2(3)(3) + 3(2)(2)} = \sqrt{30}$$

$$e) d(\vec{u}, \vec{v}) = \sqrt{\langle \vec{u} - \vec{v}, \vec{u} - \vec{v} \rangle} \quad \vec{u} - \vec{v} = \langle -2, -1 \rangle$$

$$= \sqrt{2(-2)^2 + 3(-1)^2} = \sqrt{11}$$

Theorem 6.1.1 Norms and Distances in Inner Product Spaces

If \mathbf{u} and \mathbf{v} are vectors in a real inner product space V , and if k is a scalar, then:

a) $\|\mathbf{v}\| \geq 0$ with equality if and only if $\mathbf{v} = \mathbf{0}$ (analogous to Theorem 3.2.1 (a) and (b)).

b) $\|k\mathbf{v}\| = |k| \|\mathbf{v}\|$ (analogous to Theorem 3.2.1 (c)).

c) $d(\mathbf{u}, \mathbf{v}) = d(\mathbf{v}, \mathbf{u})$ (analogous to Theorem 3.2.2 (d)).

d) $d(\mathbf{u}, \mathbf{v}) \geq 0$ with equality if and only if $\mathbf{u} = \mathbf{v}$ (ibid).

#37 Let the vector space P_2 have the inner product $\langle p, q \rangle = \int_{-1}^1 p(x)q(x)dx$. Find the following for $p = 1$ and $q = x^2$.

a. $\langle p, q \rangle$ b. $d(p, q)$ c. $\|p\|$ d. $\|q\|$

$$a) \langle \vec{p}, \vec{q} \rangle = \int_{-1}^1 x^2 dx = \frac{2}{3}$$

$$b) d(\vec{p}, \vec{q}) = \sqrt{\langle p-q, p-q \rangle} = \left[\int_{-1}^1 (1-x^2)^2 dx \right]^{1/2}$$
$$= \left(\int_{-1}^1 (1-2x^2+x^4) dx \right)^{1/2} = \frac{4}{\sqrt{15}}$$

$$c) \|\vec{p}\| = \langle \vec{p}, \vec{p} \rangle = \left(\int_{-1}^1 1 dx \right)^{1/2} = \sqrt{2}$$

$$d) \|\vec{q}\| = \langle \vec{q}, \vec{q} \rangle = \left(\int_{-1}^1 x^4 dx \right)^{1/2} = \sqrt{\frac{2}{5}}$$

Definition: If V is an inner product space, then the set of points in V that satisfy $\|u\| = 1$ is called the **unit sphere** in V (or the **unit circle** in the case where $V = \mathbb{R}^2$).

The Euclidean inner product on \mathbb{R}^2 gives us the unit circle.

A weighted Euclidean inner product gives us

a distorted circle - a.k.a. an ellipse.

In the case of the integral inner product like in # 37, the unit sphere

consists of all functions $f \in C[a, b]$

such that $\int_a^b [f(x)]^2 dx = 1$.

Ex: $\int_{-1}^1 \frac{1}{\sqrt{2}} dx = 1 \quad \Rightarrow f(x) = \frac{1}{\sqrt{2}}$

$\int_0^{\pi/2} \sqrt{\sin x} dx = 1 \quad \Rightarrow f(x) = \sqrt{\sin x}$

Theorem 6.1.2 Algebraic Properties of Inner Products (generalization of Theorem 3.2.3)

If \mathbf{u} , \mathbf{v} and \mathbf{w} are vectors in a real inner product space V , and if k is a scalar, then:

- a) $\langle \mathbf{0}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{0} \rangle = 0$
- b) $\langle \mathbf{u}, \mathbf{v} + \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle + \langle \mathbf{u}, \mathbf{w} \rangle$
- c) $\langle \mathbf{u}, \mathbf{v} - \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{v} \rangle - \langle \mathbf{u}, \mathbf{w} \rangle$
- d) $\langle \mathbf{u} - \mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{u}, \mathbf{w} \rangle - \langle \mathbf{v}, \mathbf{w} \rangle$
- e) $k \langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{u}, k\mathbf{v} \rangle$

proving these is a good exercise